

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

same part of *Jupiter's* orbit that the planet and Comet 1889 V were in July 20, 1886. The interval between these disturbances is 107 years. The period of Comet 1889 V being about 31 years, which is not an aliquot part of 107, that comet could not have been near *Jupiter* in 1779, unless it suffered other and serious disturbances in the intervening years. Such disturbances did take place in 1827 and 1791, but they were of such a nature as to leave very serious doubts whether the comet was near *Jupiter* in 1779, a condition absolutely necessary to establishing the identity of the two comets.

Dr. Poor concludes that the vexed question of identity cannot now be answered, but we must await further observations of the comet at its reappearance in 1896. He promises in another paper to discuss the question of the disruption of the comet while in *Jupiter's* system, and of the possibility of a portion of it being permanently drawn into the system to form a new satellite.

W. W. C.

The Parallax of the Planetary Nebula DM.
$$+41^{\circ},4004$$
.

Dr. J. Wilsing, of the Potsdam Astro-Physical Observatory, has determined the parallax of Webb's planetary nebula DM. + 41°, 4004 by the photographic method. From June, 1892, to June, 1893, 102 exposures of eight minutes each were made on the region of the nebula. The distances of the nebula from two eleventh magnitude stars were accurately obtained from all the plates, and a combination of the data gave as the most probable value of the parallax of the nebula relative to one of the stars

$$\pi = -0''.08$$
:

and relative to the other star

$$\pi = -0''$$
. 17.

The fact that the relative parallax comes out negative indicates that the distance of the nebula from our solar system is probably greater than that of the two eleventh magnitude comparison stars.

This is the first nebula parallax to be investigated photographically. Dr. Wilsing's paper is published in *Astronomische Nachrichten*, No. 3190. W. W. C.